

# UNIVERSAL FAMILIES ON MODULI SPACES OF PRINCIPAL BUNDLES ON CURVES

V. BALAJI, I. BISWAS, D. S. NAGARAJ, AND P. E. NEWSTEAD

ABSTRACT. Let  $H$  be a connected semisimple linear algebraic group defined over  $\mathbb{C}$  and  $X$  a compact connected Riemann surface of genus at least three. Let  $\mathcal{M}'_X(H)$  be the moduli space parametrising all topologically trivial stable principal  $H$ -bundles over  $X$  whose automorphism group coincides with the centre of  $H$ . It is a Zariski open dense subset of the moduli space of stable principal  $H$ -bundles. We prove that there is a universal principal  $H$ -bundle over  $X \times \mathcal{M}'_X(H)$  if and only if  $H$  is an adjoint group (that is, the centre of  $H$  is trivial).

## 1. INTRODUCTION

Let  $X$  be a compact connected Riemann surface of genus at least three. Let  $\mathcal{M}_X(n, d)$  denote the moduli space of all stable vector bundles over  $X$  of rank  $n$  and degree  $d$ , which is a smooth irreducible quasiprojective variety defined over  $\mathbb{C}$ . A vector bundle  $\mathcal{E}$  over  $X \times \mathcal{M}_X(n, d)$  is called *universal* if for every point  $m \in \mathcal{M}_X(n, d)$ , the restriction of  $\mathcal{E}$  to  $X \times \{m\}$  is in the isomorphism class of holomorphic vector bundles over  $X$  defined by  $m$ . A well-known theorem says that there is a universal vector bundle over  $X \times \mathcal{M}_X(n, d)$  if and only if  $d$  is coprime to  $n$  (see [13] for existence in the coprime case, [8] for non-existence in the non-coprime case and [7] for a topological version of non-existence in the case  $d = 0$ ).

Let  $H$  be a connected semisimple linear algebraic group defined over the field of complex numbers. Ramanathan extended the notion of (semi)stability to principal  $H$ -bundles and constructed moduli spaces for stable principal  $H$ -bundles over  $X$  [9, 10, 11]. The construction works for any given topological type, yielding a moduli space which is an irreducible quasiprojective variety defined over  $\mathbb{C}$ . We are concerned here with the case of topologically trivial stable principal  $H$ -bundles. Let  $\mathcal{M}_X(H)$  denote the moduli space of topologically trivial stable principal  $H$ -bundles over  $X$ .

Let  $Z(H) \subset H$  be the centre. For any  $H$ -bundle  $E_H$ , the group  $Z(H)$  is contained in the automorphism group  $\text{Aut}(E_H)$ . Let

$$\mathcal{M}'_X(H) \subset \mathcal{M}_X(H)$$

be the subvariety consisting of all  $H$ -bundles  $E_H$  over  $X$  with the property  $\text{Aut}(E_H) \cong Z(H)$ . It is known that  $\mathcal{M}'_X(H)$  is a dense Zariski-open subset contained in the smooth locus of  $\mathcal{M}_X(H)$ .

---

All authors are members of the international research group VBAC (Vector Bundles on Algebraic Curves), which is partially supported by EAGER (EC FP5 Contract no. HPRN-CT-2000-00099) and by EDGE (EC FP5 Contract no. HPRN-CT-2000-00101). The fourth author would also like to thank the Royal Society and CSIC, Madrid, for supporting a visit to CSIC during which his contribution to this work was completed.

A principal  $H$ -bundle  $E$  over  $X \times \mathcal{M}'_X(H)$  will be called a *universal bundle* if for every point  $m \in \mathcal{M}_X(n, d)$ , the restriction of  $\mathcal{E}$  to  $X \times \{m\}$  is in the isomorphism class of stable  $H$ -bundles over  $X$  defined by the point  $m$  of the moduli space.

The following theorem is the main result proved here:

**Theorem 1.1.** *There is a universal  $H$ -bundle over  $X \times \mathcal{M}'_X(H)$  if and only if  $Z(H) = e$ .*

**Remark 1.2.** Non-existence for  $H = \mathrm{SL}(n, \mathbb{C})$  was previously known [8, 7], as also was existence for  $H = \mathrm{PGL}(n, \mathbb{C})$ .

*Acknowledgments.* We thank A. King, A. Nair and P. Sankaran for useful discussions.

## 2. EXISTENCE OF UNIVERSAL BUNDLE

We begin this section by recalling very briefly certain facts from [2]. Unless otherwise specified, all bundles and sections considered will be algebraic.

**Remark 2.1.** Let  $E$  be a principal  $G$ -bundle over  $X$ , where  $G$  is a reductive linear algebraic group defined over  $\mathbb{C}$ , and  $H \subset G$  is a Zariski closed semisimple subgroup. For any variety  $Y$  equipped with an action of  $G$ , the fibre bundle  $(E \times Y)/G$  over  $X$  associated to  $E$  will be denoted by  $E(Y)$ .

- (1) There is a natural action of the group  $\mathrm{Aut}_G E$ , defined by all automorphisms of  $E$  over the identity map of  $X$  that commute with the action of  $G$ , on  $\Gamma(X, E(G/H))$  (the space of all holomorphic sections of the fibre bundle  $E(G/H) = E/H$  over  $X$ ) and the orbits correspond to the equivalence classes of  $H$ -reductions of  $E$  with two reductions being equivalent if the corresponding principal  $H$ -bundles are isomorphic.
- (2) Let  $G = \mathrm{GL}(n, \mathbb{C})$ , and let  $\phi : H \hookrightarrow G$  be a faithful representation of the semisimple group  $H$ . Let  $Q$  denote the open subset of semistable principal  $G$ -bundles (or equivalently, of trivial determinant semistable vector bundles of rank  $n$ ) of the usual “Quot scheme”, and let  $Q(\phi)$  be the “Quot scheme” which parametrises pairs of the form  $(E', s)$ , where  $E'$  is a principal  $G$ -bundle and  $s$  is a reduction of structure group of  $E'$  to  $H$ . Then  $Q(\phi)$  is in a sense a “relative Quot scheme”. As is clear from the definition and the notation, this scheme is dependent on the choice of the inclusion  $\phi : H \hookrightarrow G$  (for details see [2, 10, 11]).

One also has a *tautological* sheaf on  $X \times Q$  which in fact is a vector bundle. We denote by  $\mathcal{E}$  the *associated tautological* principal  $G$ -bundle on  $X \times Q$ .

Recall that the moduli space of principal  $G$ -bundles ( $G = \mathrm{GL}(n, \mathbb{C})$ ) is realised as a good quotient of  $Q$  by the action of a reductive group  $\mathcal{G}$ . We may also assume that the group  $\mathcal{G}$  is with trivial centre (see for example [4]).

**Remark 2.2.** It is immediate that the action of  $\mathcal{G}$  on  $Q$  lifts to an action on  $Q(\phi)$ , where  $\mathcal{G}$ ,  $Q$  and  $Q(\phi)$  are defined in Remark 2.1.

We have a morphism

$$(1) \quad \psi : Q(\phi) \longrightarrow Q,$$

which sends any  $H$ -bundle  $E'$  to the  $\mathrm{GL}(n, \mathbb{C})$ -bundle obtained by extending the structure group of  $E'$  using the homomorphism  $\phi$ . In fact,  $\psi$  is a  $\mathcal{G}$ -equivariant *affine morphism* (see [2]).

Continuing with the notation in the above two remarks, consider the  $\mathcal{G}$ -action on  $Q(\phi)$  (defined in Remark 2.1(2)) with the linearisation induced by the affine  $\mathcal{G}$ -morphism  $\psi$  in (1). Since a good quotient of  $Q$  by  $\mathcal{G}$  exists and since  $\psi$  is an affine  $\mathcal{G}$ -equivariant map, a good quotient  $Q(\phi)//\mathcal{G}$  exists (see [11, Lemma 5.1]).

Moreover by the universal property of categorical quotients, the canonical morphism

$$(2) \quad \overline{\psi} : Q(\phi)//\mathcal{G} \longrightarrow Q//\mathcal{G}$$

given by  $\psi$  is also *affine*.

**Theorem 2.3.** *Let  $\mathcal{M}_X(H)$  denote the scheme  $Q(\phi)//\mathcal{G}$  (see (2)). Then this scheme is the coarse moduli scheme of semistable  $H$ -bundles. Further, the scheme  $\mathcal{M}_X(H)$  is projective, and if  $H \hookrightarrow \mathrm{GL}(V)$  is a faithful representation, then the canonical morphism*

$$\overline{\psi} : \mathcal{M}_X(H) \longrightarrow \mathcal{M}_X(\mathrm{GL}(V)) = Q//\mathcal{G}$$

*is finite.*

Let  $Q(\phi)^s$  be the open subscheme of  $Q(\phi)$  consisting of stable  $H$ -bundles.

**Lemma 2.4.** *Let  $Q'_H \subset Q(\phi)^s$  be the subset parametrising all stable  $H$ -bundles whose automorphism group is  $Z(H)$ . Then the action of  $\mathcal{G}$  on the subset  $Q'_H$  is free, and furthermore, the quotient morphism  $Q'_H \longrightarrow Q'_H/\mathcal{G}$  is a principal  $\mathcal{G}$ -bundle. In fact,  $Q'_H/\mathcal{G}$  is precisely the Zariski open subset  $\mathcal{M}'_X(H)$  (the variety  $\mathcal{M}'_X(H)$  is defined in the introduction).*

*Proof.* For any point  $E_H \in Q(\phi)$ , the isotropy subgroup of  $E_H$  for the action of  $\mathcal{G}$  coincides with  $\mathrm{Aut}(E_H)/Z(H)$ . This can be seen as follows: firstly, the point  $E_H$  is a pair  $(E, s)$ , where  $E \in Q$  and  $s$  is a reduction of structure group to  $H$  of the  $G$ -bundle  $E$ . It is well known for the action of  $\mathcal{G}$  on  $Q$  that the isotropy at  $E$  is precisely the group  $\mathrm{Aut}(E)/Z(G)$ . From this it is easy to see that the isotropy of  $E_H$  for the action of  $\mathcal{G}$  on  $Q(\phi)$  is the group  $\mathrm{Aut}(E, s)/Z(H) = \mathrm{Aut}(E_H)/Z(H)$ .

Hence it follows that the action of  $\mathcal{G}$  on the open subset  $Q'_H$  is free, and the proof of the lemma is complete.  $\square$

We remark that, at least when  $H$  is of adjoint type,  $Q'_H$  is a non-empty open subset of  $Q(\phi)^s$ . Openness is easy and can be seen for example from [5, Theorem II.6 (ii)]. Non-emptiness follows from Proposition 2.6 below.

We prove a proposition on semisimple groups, possibly known to experts but which we could not locate in any standard text.

**Proposition 2.5.** *Let  $H$  be a semisimple algebraic group. Then  $H$  has a faithful, irreducible representation  $\phi : H \longrightarrow \mathrm{GL}(V)$  if and only if the centre of  $H$  is cyclic.*

*Proof.* One way the implication is easy, namely suppose that a faithful irreducible representation exists, then the centre  $Z(H)$  of  $H$  is cyclic. To see this, first note that under the representation  $\phi$ , the centre  $Z(H)$  maps to a subgroup which commutes with all elements of  $\phi(H)$ . Since  $\phi$  is irreducible, this implies by Schur's Lemma that  $\phi(Z(H)) \subset Z(\mathrm{GL}(V))$ , where  $Z(\mathrm{GL}(V))$  is the centre of  $\mathrm{GL}(V)$ . Observe further that since  $H$  is semisimple, we have  $\phi(H) \subset \mathrm{SL}(V)$ . Hence,  $Z(H) \subset Z(\mathrm{SL}(V))$  and is therefore cyclic.

The contrapositive statement is harder to prove. We proceed as follows: Let  $H'$  be the simply connected cover of  $H$ , and let  $\overline{H}$  be the associated adjoint group, namely  $H/Z(H)$ . Let  $\Lambda$  (respectively,  $\Lambda_R$ ) be the weight lattice (respectively, root lattice). In other words, by the Borel-Weil theorem  $\Lambda = \mathcal{X}(B)$  and  $\Lambda_R = \mathcal{X}(\overline{B})$ , where  $B, \overline{B}$  are fixed Borel subgroups of  $H, \overline{H}$ . Then, from the exact sequence

$$e \longrightarrow Z(H) \longrightarrow H \longrightarrow \overline{H} \longrightarrow e$$

we see that  $\Lambda/\Lambda_R \simeq \mathcal{X}(Z(H))$ . In other words, the quotient group  $\Lambda/\Lambda_R$  is cyclic of order  $m$ .

Let  $\overline{\lambda}$  be a generator of the cyclic group  $\Lambda/\Lambda_R$ . Then, by an action of the Weyl group we may assume that the coset representative  $\lambda \in \Lambda$  is actually a dominant weight.

Suppose that the root lattice has the following decomposition (corresponding to the simple components of  $\overline{H}$ ):

$$\Lambda_R = \bigoplus_{i=1}^{\ell} \Lambda^i.$$

Then, by possibly adding dominant weights from the  $\Lambda^i$ , we may assume that  $(m \cdot \lambda) \in \Lambda_R$  has all its direct sum components  $\lambda_i \neq 0$ ,  $i \in [1, \ell]$ , where  $\lambda$  is as above.

For this choice of  $\lambda \in \Lambda$  let  $V_\lambda$  be the corresponding  $H$ -module given by

$$\phi_\lambda : H \longrightarrow \mathrm{GL}(V_\lambda).$$

Then one knows that  $\phi_\lambda$  is an irreducible representation of  $H$ .

We claim that the representation  $\phi_\lambda$  is even *faithful*. Suppose that this is not the case. Let  $K_\lambda := \ker(\phi_\lambda) \neq e$ .

Firstly,  $K_\lambda \subset Z(H)$ . To see this, let  $K'$  be the inverse image of  $K_\lambda$  in the simply connected cover  $H'$  of  $H$ . Then the choice of  $\lambda$  so made that its simple components are non-zero in fact forces the following: Suppose that  $H' = H_1 \times \cdots \times H_\ell$  is the decomposition of  $H'$  into its almost simple factors. (We recall that a semisimple algebraic group is called almost simple if the quotient of it by its centre is simple.) Then the normal subgroup  $K'$  in its decomposition in  $H'$  is such that  $K_i := K' \cap H_i$  are proper normal subgroups of  $H_i$ . In particular,  $K_i \subset Z(H_i)$  for all  $i \in [1, \ell]$ . This implies that  $K' \subset Z(H')$  and hence,  $K_\lambda \subset Z(H)$ .

Note that the dominant character  $\lambda$  is non-trivial on the generator of the centre  $Z(H)$  because  $\mathcal{X}(Z(H)) = \Lambda/\Lambda_R$ . Now  $K_\lambda \subset Z(H)$  and  $Z(H)$  cyclic implies that  $\lambda$  is non-trivial on the generator of  $K_\lambda$  as well. This contradicts the fact that  $K_\lambda = \ker(\phi_\lambda)$ . This proves the claim. Therefore, the proof of the proposition is complete.  $\square$

If  $H$  is of adjoint type, then by Proposition 2.5 we can choose the inclusion  $\phi : H \hookrightarrow G$  in Remark 2.1(2) to be an irreducible representation. Henceforth,  $\phi$  will be assumed to be irreducible.

Let  $E$  be a stable  $H$ -bundle of trivial topological type. Recall that one can realise  $E$  from a unique, up to an inner conjugation, irreducible representation of  $\pi_1(X)$  in a maximal compact subgroup of  $H$  (see [9]). For notational convenience, we will always suppress the base point in the notation of fundamental group. Denote by  $M(E)$  the Zariski closure of the image of  $\pi_1(X)$  in  $H$ .

**Proposition 2.6.** *Let  $H$  be of adjoint type and let  $\phi$  be a faithful irreducible representation of  $H$  in  $V$  (see Proposition 2.5). Let  $U \subset Q(\phi)^s$  be the subset defined as follows:*

$$U = \{E \in Q(\phi)^s \mid M(E) = H\}.$$

*Then  $U$  is non-empty and is contained in the subset  $Q'_H$  of stable  $H$ -bundles which have trivial automorphism group. (Since  $H$  is of adjoint type,  $Z(H) = e$ .)*

*Proof.* Since  $\phi$  is irreducible, using Lemma 2.1 of [12] we conclude that there is an irreducible representation

$$\rho : \pi_1(X) \longrightarrow H$$

which have the property that the composition  $\phi \circ \rho : \pi_1(X) \longrightarrow G := \mathrm{GL}(V)$  continues to remain irreducible. This implies that  $M(E_\rho) = H$  by the construction in [12, Lemma 2.1] and hence  $E_\rho \in U$ , i.e.,  $U$  is non-empty.

Let  $E \in U$ . Then by the definition of  $U$ , there exists a representation  $\rho$  of  $\pi_1(X)$  in  $H$  such that  $E \simeq E_\rho$ , where  $E_\rho$  is the flat principal  $H$ -bundle given by  $\rho$ . Observe that  $E_\rho$  is a stable  $H$ -bundle and the associated  $G$ -bundle is also stable. Hence all the automorphisms of this associated  $G$ -bundle lie in  $Z(G)$ . Since  $H$  is of adjoint type, it follows that the  $H$ -bundle  $E_\rho$  has no non-trivial automorphisms. Hence it follows that  $U \subset Q'_H$ .  $\square$

We have the following theorem on existence of universal families.

**Theorem 2.7.** *Let  $H$  be a group of adjoint type and  $\mathcal{M}'_X(H)$  the Zariski open subset of  $\mathcal{M}_X(H)$  defined in the introduction. Then there exists a universal family of principal  $H$ -bundles on  $X \times \mathcal{M}'_X(H)$ .*

*Proof.* We first observe that the variety  $\mathcal{M}'_X(H)$  is precisely the image of  $Q'_H$  under the quotient map for the action of  $\mathcal{G}$ . We recall that  $Q'_H$  is nonempty by Proposition 2.6.

Since  $Z(H) = e$ , it follows that  $H \subset \overline{G} := G/Z(G)$ , where  $G = \mathrm{GL}(V)$  is as in Remark 2.1(2).

Consider the tautological  $G$ -bundle  $\mathcal{E}$  on  $X \times Q^s$ , and let  $\overline{\mathcal{E}}$  be the corresponding  $\overline{G}$ -bundle obtained by extending the structure group. Then it is well-known that the adjoint universal bundle  $\overline{\mathcal{E}}$  descends to the quotient  $\mathcal{M}_X(G)^s$  (see [4]).

We follow the same strategy for  $\mathcal{M}'_X(H)$  as well. Consider the pulled back  $\overline{G}$ -bundle  $(\mathrm{Id}_X \times \psi)^*\overline{\mathcal{E}}$  on  $X \times Q'_H$ , where  $\psi$  is the map in (1) and  $\overline{G} := G/Z(G)$ . The action of  $\mathcal{G}$  on  $Q'_H$  is free by Lemma 2.4. Therefore, the quotient  $Q'_H \longrightarrow \mathcal{M}'_X(H)$  is a principal  $\mathcal{G}$ -bundle. Further, the action of  $\mathcal{G}$  lifts to the tautological bundle  $(\mathrm{Id}_X \times \psi)^*\overline{\mathcal{E}}$ . In particular, the principal  $\overline{G}$ -bundle  $(\mathrm{Id}_X \times \psi)^*\overline{\mathcal{E}}$  descends to a principal  $\overline{G}$ -bundle over  $X \times \mathcal{M}'_X(H)$ .

Let us denote this descended  $\overline{G}$ -bundle over  $X \times \mathcal{M}'_X(H)$  by  $\overline{\mathcal{E}}_0$ .

Let  $\pi : (\mathrm{Id}_X \times \psi)^*\mathcal{E}(G/H) \longrightarrow (\mathrm{Id}_X \times \psi)^*\overline{\mathcal{E}}(\overline{G}/H)$  be the natural map induced by the projection  $G/H \longrightarrow \overline{G}/H$ , where  $\psi$  is the map in (1). We note that the universal  $H$ -bundle over  $X \times U$ , where  $U$  is defined in Proposition 2.6, is a reduction of structure group of the pulled back  $G$ -bundle  $(\mathrm{Id}_X \times \psi)^*\mathcal{E}$ . Let

$$\sigma : X \times U \longrightarrow (\mathrm{Id}_X \times \psi)^*\mathcal{E}(G/H)$$

be the section giving this reduction of structure group. Then the composition  $\pi \circ \sigma$  is a section of  $(\text{Id}_X \times \psi)^* \bar{\mathcal{E}}(\bar{G}/H)$  over  $X \times U$ .

Since  $H$  is semisimple, a lemma of Chevalley says that there is a  $\bar{G}$ -module  $W$  and an element  $w \in W$  such that  $H$  is precisely the isotropy subgroup for  $w$  (see [3, p. 89, Theorem 5.1]). Therefore,  $\bar{G}/H$  is identified with the closed  $\bar{G}$ -orbit in  $W$  defined by  $w$ . Then we see that  $\bar{\mathcal{E}}(\bar{G}/H) \hookrightarrow \bar{\mathcal{E}}(W)$ . We may therefore view the section  $\pi \circ \sigma : X \times U \longrightarrow (\text{Id}_X \times \psi)^* \bar{\mathcal{E}}(\bar{G}/H)$  as a section of the vector bundle  $(\text{Id}_X \times \psi)^* \bar{\mathcal{E}}(W)$  over  $X \times U$ .

Since  $(\text{Id}_X \times \psi)^* \bar{\mathcal{E}}$  descends to the  $\bar{G}$ -bundle  $\bar{\mathcal{E}}_0$  over  $X \times \mathcal{M}'_X(H)$ , it follows that the associated vector bundle  $(\text{Id}_X \times \psi)^* \bar{\mathcal{E}}(W)$  also descends to  $X \times \mathcal{M}'_X(H)$ . Clearly this vector bundle is nothing but the associated vector bundle  $\bar{\mathcal{E}}_0(W)$ , associated to the  $\bar{G}$ -bundle  $\bar{\mathcal{E}}_0$  on  $X \times \mathcal{M}'_X(H)$  for the  $\bar{G}$ -module  $W$ .

Since each point of  $\mathcal{M}'_X(H)$  represents an isomorphism class of stable  $H$ -bundle, it follows that set theoretically the reduction section  $\pi \circ \sigma \in \Gamma((\text{Id}_X \times \psi)^* \bar{\mathcal{E}}(W))$  descends to a section on  $X \times \mathcal{M}'_X(H)$  of the descended vector bundle  $\bar{\mathcal{E}}_0(W)$ .

We now appeal to [6, Proposition 4.1], which implies that the section  $\pi \circ \sigma$  in fact descends to give a holomorphic section of  $\bar{\mathcal{E}}_0(W)$  over  $X \times \mathcal{M}'_X(H)$ .

Again set-theoretically, the image of this section of  $\bar{\mathcal{E}}_0(W)$  lies in  $\bar{\mathcal{E}}_0(\bar{G}/H) \subset \bar{\mathcal{E}}_0(W)$ . As before, from [6, Proposition 4.1] it follows that  $\pi \circ \sigma$  gives a reduction of structure group to  $H$  of the descended  $\bar{G}$ -bundle  $\bar{\mathcal{E}}_0$ . The  $H$ -bundle over  $X \times \mathcal{M}'_X(H)$  obtained this way is the required universal  $H$ -bundle. This completes the proof of the theorem.  $\square$

### 3. NONEXISTENCE OF UNIVERSAL BUNDLE

Let  $H$  be a complex semisimple linear algebraic group, and let  $K \subset H$  be a maximal compact subgroup. The Lie algebra of  $K$  will be denoted by  $\mathfrak{k}$ . A homomorphism

$$\rho : \pi_1(X) \longrightarrow K$$

is called *irreducible* if no nonzero vector in  $\mathfrak{k}$  is fixed by the adjoint action of the subgroup  $\rho(\pi_1(X)) \subset K$  on  $\mathfrak{k}$ . Let  $\text{Hom}^{\text{irr}}(\pi_1(X), K)$  denote the space of all irreducible homomorphisms such that the corresponding  $K$ -bundle is topologically trivial. So any homomorphism in  $\text{Hom}^{\text{irr}}(\pi_1(X), K)$  is induced by a homomorphism from  $\pi_1(X)$  to the universal cover of  $K$ .

Let  $g$  denote the genus of  $X$ . Assume that  $g \geq 3$ . If we choose a basis

$$\{a_1, \dots, a_g, b_1, \dots, b_g\} \subset \pi_1(X)$$

such that

$$(3) \quad \pi_1(X) = \langle a_1, \dots, a_g, b_1, \dots, b_g : \prod_{i=1}^n a_i b_i a_i^{-1} b_i^{-1} \rangle,$$

then  $\text{Hom}^{\text{irr}}(\pi_1(X), K)$  gets identified with a real analytic subspace of  $K^{2g}$ .

For any  $\rho \in \text{Hom}^{\text{irr}}(\pi_1(X), K)$ , the principal  $H$ -bundle obtained by extending the structure group of the principal  $K$ -bundle given by  $\rho$  is stable. A theorem of Ramanathan says that all topologically trivial stable  $H$ -bundles arise in this way, that is, the space of all equivalence classes of irreducible homomorphisms of  $\pi_1(X)$  to  $K$  is in bijective correspondence with the space of all stable  $H$ -bundles over  $X$  [9, Theorem 7.1]. More precisely

(see the proof of [9, Theorem 7.1]), the real analytic space underlying the moduli space  $\mathcal{M}_X(H)$  parametrising topologically trivial stable  $H$ -bundles is analytically isomorphic to the quotient space

$$R(\pi_1(X), H) := \text{Hom}^{\text{irr}}(\pi_1(X), K)/K,$$

for the action constructed using the conjugation action of  $K$  on itself. Let

$$(4) \quad q : \text{Hom}^{\text{irr}}(\pi_1(X), K) \longrightarrow R(\pi_1(X), H) \cong \mathcal{M}_X(H)$$

be the quotient map. The open subset  $\mathcal{M}'_X(H) \subset \mathcal{M}_X(H)$  is a smooth submanifold and the restriction of the map in (4)

$$(5) \quad q|_{q^{-1}(\mathcal{M}'_X(H))} : q^{-1}(\mathcal{M}'_X(H)) \longrightarrow \mathcal{M}'_X(H)$$

is a smooth principal  $K/Z$ -bundle, where  $Z$  is the centre of  $K$ . Note that  $Z$  coincides with the centre of  $H$  (as  $H$  is semisimple, its centre is a finite group).

If  $\mathcal{U}_H$  is a universal  $H$ -bundle over  $X \times \mathcal{M}'_X(H)$ , then we have a reduction of structure group of  $\mathcal{U}_H$  to  $K$  which is constructed using the correspondence established in [9] between stable  $H$ -bundles and irreducible flat  $K$ -bundles. Let

$$(6) \quad \mathcal{U}_K \longrightarrow X \times \mathcal{M}'_X(H)$$

be the smooth principal  $K$ -bundle obtained from  $\mathcal{U}_H$  this way. Consider the principal  $K/Z$ -bundle over  $X \times \mathcal{M}'_X(H)$ , where  $Z \subset K$  is the centre, obtained by extending the structure group of  $\mathcal{U}_K$  using the natural projection of  $K$  to  $K/Z$ . The restriction of this  $K/Z$ -bundle to  $p \times \mathcal{M}'_X(H)$ , where  $p$  is the base point in  $X$  used for defining  $\pi_1(X)$ , is identified with the  $K/Z$ -bundle in (5). To see this first note that the universal  $K$ -bundle over  $X \times \text{Hom}(\pi_1(X), K)$  is obtained as a quotient by the action of  $\pi_1(X)$  on  $\tilde{X} \times \text{Hom}(\pi_1(X), K) \times K$ , where  $\tilde{X}$  is the pointed universal cover of  $X$  for the base point  $p$ ; the action of  $z \in \pi_1(X)$  sends any  $(\alpha, \beta, \gamma)$  to  $(\alpha, z, \beta, \beta(z)^{-1}\gamma)$ . From this it follows that the restriction of this universal bundle to  $p \times \text{Hom}(\pi_1(X), K)$  is canonically trivialized. The above mentioned identification is constructed using this trivialization.

Our aim is to show that no  $K$ -bundle over  $\mathcal{M}'_X(H)$  exists that produces the  $K/Z$ -bundle in (5) by extension of structure group, provided the centre  $Z$  is non-trivial.

Let  $F_3$  denote the free group on three generators. Fix a surjective homomorphism

$$(7) \quad f : \pi_1(X) \longrightarrow F_3$$

that sends  $a_i$ ,  $1 \leq i \leq 3$ , to the  $i$ -th generator of  $F_3$  and sends  $a_i$ ,  $4 \leq i \leq g$ , and  $b_i$ ,  $1 \leq i \leq g$ , to the identity element, where  $a_j$ ,  $b_j$  are as in (3) (recall that  $g \geq 3$ ).

Set

$$(8) \quad R(F_3, H) := \text{Hom}^{\text{irr}}(F_3, K)/K$$

to be the equivalence classes of irreducible representations. Note that  $\text{Hom}(F_3, K) \simeq K^3$ , and under this identification the action

$$\mu : \text{Hom}(F_3, K) \times K \longrightarrow \text{Hom}(F_3, K)$$

given by  $\mu(\rho, A) = A^{-1}\rho A$  corresponds to the simultaneous diagonal conjugation action of  $K$  on the three factors.

Since  $F_3$  is a free group and  $K$  is connected, any homomorphism  $\rho$  from  $F_3$  to  $K$  can be deformed to the trivial homomorphism. This implies that the principal  $H$ -bundle corresponding to the homomorphism

$$\rho \circ f : \pi_1(X) \longrightarrow K$$

is topologically trivial, where  $f$  is defined in (7). Therefore, we have an embedding

$$(9) \quad R(F_3, H) \longrightarrow R(\pi_1(X), H) \cong \mathcal{M}_X(H)$$

that sends any  $\rho$  to  $\rho \circ f$ . Let  $R'(F_3, H) \subset R(F_3, H)$  be the inverse image of the open subset  $\mathcal{M}'_X(H)$  under the above map. By Proposition 2.6 it follows that  $R'(F_3, H)$  is a *non-empty* open subset of  $R(F_3, H)$ .

Fix a point  $p \in X$ , which will also be the base point for the fundamental group. Consider the restriction of the principal  $K$ -bundle  $\mathcal{U}_K$  in (6) to  $p \times \mathcal{M}'_X(H) \hookrightarrow X \times \mathcal{M}'_X(H)$  and denote it by  $\mathcal{U}_{K,p}$ .

Let

$$(10) \quad \gamma : p \times R'(F_3, H) \longrightarrow p \times \mathcal{M}'_X(H)$$

be the map given by the embedding  $R'(F_3, H) \longrightarrow \mathcal{M}'_X(H)$  constructed in (9). Taking the pull back of the above defined principal  $K$ -bundle  $\mathcal{U}_{K,p}$  over  $p \times \mathcal{M}'_X(H)$  under the morphism  $\gamma$  in (10),

$$(11) \quad \begin{array}{ccc} \widetilde{M} & & \mathcal{U}_{K,p} \\ \downarrow & & \downarrow \\ p \times R'(F_3, H) & \xrightarrow{\gamma} & p \times \mathcal{M}'_X(H) \end{array}$$

we obtain a principal  $K$ -bundle  $\widetilde{M} := \gamma^* \mathcal{U}_{K,p}$  on  $R'(F_3, H)$  whose associated  $K/Z$ -bundle (as before,  $Z$  is the centre of  $K$ ) is precisely the  $K/Z$ -bundle  $q^{-1}(R'(F_3, H)) \longrightarrow R'(F_3, H)$  constructed in (5). In particular,  $q^{-1}(R'(F_3, H)) \simeq \widetilde{M}/Z$ .

Let  $\text{Hom}^{\text{irr}}(F_3, K)' \subset \text{Hom}^{\text{irr}}(F_3, K)$  be the inverse image of  $R'(F_3, H)$  under the quotient map in (8). Let

$$(12) \quad q_0 : \text{Hom}^{\text{irr}}(F_3, K)' \longrightarrow R'(F_3, H)$$

be the restriction of the quotient map in (8). So  $q_0$  is the quotient for the conjugation action of  $K$ . This map  $q_0$  defines a principal  $K/Z$ -bundle over  $R'(F_3, H)$  which is evidently identified with the  $K/Z$ -bundle  $\widetilde{M}/Z \longrightarrow R'(F_3, H)$  obtained from (11).

We will prove (by contradiction) that such a  $K$ -bundle  $\widetilde{M}$  does not exist. In other words, we will show that there is no principal  $K$ -bundle over  $R'(F_3, H)$  whose extension of structure group is the  $K/Z$ -bundle in (12).

Assume the contrary, then we get the following diagram of topological spaces and morphisms:

$$(13) \quad \begin{array}{ccccc} Z & \longrightarrow & K & \longrightarrow & K/Z \\ || & & \downarrow & & \downarrow \\ Z & \longrightarrow & \widetilde{M} & \longrightarrow & q_0^{-1}(R'(F_3, H)) \\ & & \downarrow & & \downarrow \\ & & R'(F_3, H) & = & R'(F_3, H) \end{array}$$

We will need a couple of results. The proof of the non-existence of the bundle  $\widetilde{M}$  will be completed after establishing Proposition 3.2.

**Lemma 3.1.** *There exist finitely many compact differentiable manifolds  $N_i$  and differentiable maps*

$$f_i : N_i \longrightarrow K^3,$$

$1 \leq i \leq d$ , such that if  $N := \bigcup_{i=1}^d f_i(N_i)$  then

- (1) *the complement  $K^3 \setminus N$  is contained in  $q_0^{-1}(R'(F_3, H))$ , where  $q_0$  is the projection in (12).*
- (2) *Furthermore,  $\dim K^3 - \dim N_i \geq 4$  for all  $i \in [1, d]$ .*

*Proof.* Take any homomorphism  $\rho : \pi_1(X) \longrightarrow K$ . Let  $E_\rho$  denote the corresponding polystable principal  $H$ -bundle over  $X$  [9]. The automorphism group of the polystable  $H$ -bundle  $E_\rho$  coincides with the centraliser of  $\rho(\pi_1(X))$  in  $H$ . Hence if  $\rho(\pi_1(X))$  is dense in  $K$ , then the automorphism group of  $E_\rho$  is the centre  $Z(H) \subset H$ . Therefore, if the topological closure  $\overline{\rho(\pi_1(X))}$  of  $\rho(\pi_1(X))$  is  $K$ , and  $E_\rho$  is topologically trivial, then  $E_\rho \in \mathcal{M}'_X(H)$ .

Now assume that the centraliser  $C(\rho) \subset H$  of  $\rho(\pi_1(X))$  in  $H$  properly contains  $Z(H)$ . Since the complexification of  $\overline{\rho(\pi_1(X))}$  is reductive, and the centraliser of a reductive group is reductive, we conclude that  $C(\rho)$  is reductive. Take a semisimple element  $z \in (H \setminus Z(H)) \cap C(\rho)$  (since  $C(\rho)$  is reductive and larger than  $Z(H)$  such an element exists). Let  $C_z \subset K$  be the centraliser of  $z$  in  $K$ . We have  $\overline{\rho(\pi_1(X))} \subset C_z$ , and  $C_z$  is a proper subgroup of  $K$  as  $z \notin Z(H)$ . Also,  $C_z$  contains a maximal torus of  $K$ .

Fix a maximal torus  $T \subset K$ . (Since any two maximal tori are conjugate, any subgroup  $C_z$  of the above type would contain  $T$  after an inner conjugation.) Consider all proper Lie subgroups  $M$  of  $K$  satisfying the following two conditions:

- (i)  $T \subseteq M$ , and
- (ii) there exists a semisimple element  $z \in H \setminus Z(H)$  such that  $M$  is the centraliser of  $z$  in  $K$ .

Let  $\mathcal{S}$  denote this collection of Lie subgroups of  $K$ .

The connected ones among  $\mathcal{S}$  are precisely the maximal compact subgroups of Levi subgroups of proper parabolic subgroups of  $H$  containing  $T$ . Note that if  $P$  is a proper parabolic subgroup of  $H$ , then  $P \cap K$  is a maximal compact subgroup of a Levi subgroup of  $P$ . All the connected ones among  $\mathcal{S}$  arise as  $P \cap K$  for some proper parabolic subgroup  $P \subset H$  containing  $T$ .

This collection  $\mathcal{S}$  is a finite set. To see this, we first note that there are only finitely many parabolic subgroups of  $H$  that contain  $T$ . For a proper parabolic subgroup  $P \subset H$  containing  $T$ , there are only finitely many Lie subgroups of  $K$  that have  $P \cap K$  as the connected component containing the identity element. Thus  $\mathcal{S}$  is a finite set.

Let  $M_1, \dots, M_d$  be the subgroups of  $K$  that occur in  $\mathcal{S}$ .

We will show that the codimension of each  $M_i$  in  $K$  is at least two. It suffices show that for a maximal proper parabolic subgroup  $P$  of  $G$  containing  $T$ , the codimension of  $M := P \cap K$  in  $K$  is at least two.

To prove that the codimension of  $M = P \cap K$  in  $K$  is at least two, let  $\mathfrak{k}$  be the Lie algebra of  $K$ , and let  $\mathfrak{h}$  be the Lie algebra of  $T$  (the Cartan subalgebra). Let

$$\mathfrak{k} = \mathfrak{h} + \sum_{\alpha \text{ a root}} \mathfrak{k}^\alpha$$

be the root space decomposition of  $\mathfrak{k}$ . Let  $\mathfrak{m}$  be a Lie algebra of  $M$  which is a Lie subalgebra of  $\mathfrak{k}$  containing  $\mathfrak{h}$ . Then we have the decomposition

$$\mathfrak{m} = \mathfrak{h} + \sum_{\alpha \text{ a root}} \mathfrak{m}^\alpha$$

for  $\mathfrak{m}$  where each  $\mathfrak{m}^\alpha$  is irreducible for  $\mathfrak{h}$ , and hence has to coincide with one of the  $\mathfrak{k}^\alpha$ . Then  $\text{codim}_{\mathfrak{k}}(\mathfrak{m}) \geq 2$  since if  $\alpha$  is a root for the subalgebra  $\mathfrak{m}$  then so is  $-\alpha$  (see [1, p. 83, Corollary 4.15]).

Consequently, the codimension of  $M$  in  $K$  is at least two. Thus the codimension of each  $M_i$ ,  $i \in [1, d]$ , in  $K$  is at least two.

For each  $i \in [1, d]$ , let

$$(14) \quad \overline{f}_i : K \times M_i^3 \longrightarrow K^3$$

be the map defined by  $(x, (y_1, y_2, y_3)) \longmapsto (xy_1x^{-1}, xy_2x^{-1}, xy_3x^{-1})$ . Consider the free action of  $M_i$  on  $K \times M_i^3$  defined by

$$z \cdot (x, (y_1, y_2, y_3)) = (xz^{-1}, (zy_1z^{-1}, zy_2z^{-1}, zy_3z^{-1})),$$

where  $x \in K$  and  $z, y_1, y_2, y_3 \in M_i$ . The map  $f_i$  in (14) clearly factors through the quotient

$$\frac{K \times M_i^t}{M_i}$$

for the above action. Therefore, we have

$$(15) \quad f_i : N_i := \frac{K \times M_i^t}{M_i} \longrightarrow K^t$$

induced by  $f_i$ .

To prove part (1) of the lemma, we recall the earlier remark that for any homomorphism  $\rho' \in q_0^{-1}(R'(F_3, H))$  (the map  $q_0$  is defined in (12)), the automorphism group of the principal  $H$ -bundle corresponding to  $\rho' \circ f$  (the homomorphism  $f$  is defined in (7)) coincides with  $Z(H)$  if the image of  $\rho'$  is dense in  $K$ . From the properties of the collection

$\{M_1, \dots, M_d\}$  we conclude that

$$N := \bigcup_{i=1}^d f_i(N_i) \supseteq q_0^{-1}(R'(F_3, H))^c,$$

where  $f_i$  are defined in (15) and

$$q_0^{-1}(R'(F_3, H))^c \subset K^3$$

is the complement of  $q_0^{-1}(R'(F_3, H))$  in  $K^3$ .

Therefore, proof of part (1) is complete. To prove part (2), we note that

$$\dim N_i = \dim K + 2 \cdot \dim M_i$$

for all  $i \in [1, d]$ . It was shown earlier that  $\dim M_i \leq \dim K - 2$ . Therefore,

$$\dim K^3 - \dim N_i = 3 \cdot \dim K - \dim N_i \geq 3 \cdot \dim K - 3 \cdot \dim K + 4 = 4.$$

This completes the proof of the lemma.  $\square$

**Proposition 3.2.** *Consider the  $K/Z$ -principal bundle  $q_0^{-1}(R'(F_3, H)) \longrightarrow R'(F_3, H)$ , where  $q_0$  is the projection in (12). The induced homomorphism on fundamental groups*

$$\pi_1(K/Z) \longrightarrow \pi_1(q^{-1}(R'(F_3, H)))$$

*obtained from the homotopy exact sequence is trivial.*

*Proof.* Let  $x_1, x_2, x_3 \in K$  be regular elements; we recall that  $x \in K$  is called regular if the centralizer  $C(x) = \{y \in K \mid yx = xy\}$  is a maximal torus in  $K$ . Since the set of regular elements is dense in  $K$ , and  $q_0^{-1}(R'(F_3, H)) \subset K^3$  is a nonempty open dense subset (this follows from Lemma 3.1), we may choose these  $x_i$ ,  $i = 1, 2, 3$ , to lie in  $q_0^{-1}(R'(F_3, H))$ .

Consider the orbit  $\text{Orb}_{K^3}(x_1, x_2, x_3)$ , of  $(x_1, x_2, x_3) \in K^3$  for the adjoint action of the group  $K^3$  on itself. Clearly we have the following identification of this orbit:

$$\text{Orb}_{K^3}(x_1, x_2, x_3) = (K/C(x_1)) \times (K/C(x_2)) \times (K/C(x_3))$$

with  $C(x_i) \subset K$  being the centralizer of  $x_i$ . The conjugation action of  $K$  on  $\text{Hom}(F_3, K)$  coincides with the restriction of above action of  $K^3$  to the image of the diagonal map  $K \hookrightarrow K^3$ . Therefore, the fibre  $K/Z$  through the point  $(x_1, x_2, x_3) \in q_0^{-1}(R'(F_3, H))$  of the  $K/Z$ -bundle

$$q_0^{-1}(R'(F_3, H)) \longrightarrow R'(F_3, H)$$

is contained in the orbit

$$(K/C(x_1)) \times (K/C(x_2)) \times (K/C(x_3)).$$

Now if we choose a point  $(x_1, x_2, x_3) \in K^3$  which is general enough, then by the definition of the inverse image  $q_0^{-1}(R'(F_3, H))$  and Lemma 3.1(2) it follows immediately that the complement of the open dense subset

$$q_0^{-1}(R'(F_3, H)) \cap \left( \frac{K}{C(x_1)} \times \frac{K}{C(x_2)} \times \frac{K}{C(x_3)} \right) \subset \frac{K}{C(x_1)} \times \frac{K}{C(x_2)} \times \frac{K}{C(x_3)}$$

is of codimension at least four.

Since the image of  $K/Z$  in  $q_0^{-1}(R'(F_3, H))$  lies in  $(q^{-1}(R'(F_3, H))) \cap \text{Orb}_{K^3}(x_1, x_2, x_3)$ , whose complement is of codimension at least four in

$$\text{Orb}_{K^3}(x_1, x_2, x_3) = (K/C(x_1)) \times (K/C(x_2)) \times (K/C(x_3))$$

it follows that the homomorphism  $\pi_1(K/Z) \longrightarrow \pi_1(q_0^{-1}(R'(F_3, H)))$  in the proposition factors through

$$\begin{aligned} & \pi_1((q_0^{-1}(R'(F_3, H))) \cap \text{Orb}_{K^3}(x_1, x_2, x_3)) \\ &= \pi_1((K/C(x_1)) \times (K/C(x_2)) \times (K/C(x_3))) = \pi_1(K/T)^3 \end{aligned}$$

where  $T$  is a maximal torus of  $K$ .

For any maximal torus  $T \subset K$ , the quotient  $K/T$  is diffeomorphic to  $H/B$ , where  $B$  is a Borel subgroup of  $H$ . Since  $H/B$  is simply connected, we conclude that  $(K/C(x_1)) \times (K/C(x_2)) \times (K/C(x_3))$  is simply connected. This completes the proof of the proposition.  $\square$

We now complete the proof of the non-existence of the covering  $\widetilde{M} \longrightarrow q^{-1}(R'(F_3, H))$  as in (13).

Since the homomorphism of fundamental groups  $\pi_1(K/Z) \longrightarrow \pi_1(q_0^{-1}(R'(F_3, H)))$  is trivial (see Proposition 3.2), the induced covering  $K \longrightarrow K/Z$  is trivial (see the diagram (13)). But this is a contradiction to the fact that  $K$  is connected.

Therefore, we have proved the following theorem:

**Theorem 3.3.** *If the centre  $Z(H)$  is non-trivial, then there is no universal bundle over  $X \times \mathcal{M}'_X(H)$ .*

## REFERENCES

- [1] J. F. Adams, *Lectures on Lie Groups*, W. A. Benjamin, Inc, Amsterdam, 1969.
- [2] V. Balaji and C. S. Seshadri, *Semistable Principal Bundles-I (in char 0)*, Jour. Alg. **258** (2002), 321–347.
- [3] A. Borel, *Linear algebraic groups*, Second edition, Graduate Texts in Mathematics, 126, Springer–Verlag, New York, 1991.
- [4] J.-M. Drézet and M. S. Narasimhan, *Groupes de Picard des variétés de modules de fibrés semi-stables sur les courbes algébriques*, Invent. Math. **97** (1989), 53–94.
- [5] G. Faltings, *Stable  $G$ -bundles and projective connections*, Jour. Alg. Geom. **2** (1993), 507–568.
- [6] S. Kumar, M. S. Narasimhan and A. Ramanathan, *Infinite Grassmannians and moduli spaces of  $G$ -bundles*, Math. Ann. **300** (1994), 41–75.
- [7] P. E. Newstead, *A non-existence theorem for families of stable bundles*, Jour. Lond. Math. Soc. **6** (1973), 259–266.
- [8] S. Ramanan, *The moduli spaces of vector bundles over an algebraic curve*, Math. Ann. **200** (1973), 69–84.
- [9] A. Ramanathan, *Stable principal bundles on a compact Riemann surface*, Math. Ann. **213** (1975), 129–152.
- [10] A. Ramanathan, *Moduli of principal bundles over algebraic curves: I*, Proc. Ind. Acad. Sci. (Math. Sci.) **106** (1996), 301–328.
- [11] A. Ramanathan, *Moduli of principal bundles over algebraic curves: II*, Proc. Ind. Acad. Sci. (Math. Sci.) **106** (1996), 421–449.
- [12] S. Subramanian, *Mumford's example and a general construction*, Proc. Ind. Acad. Sci. (Math. Sci.) **99** (1989), 197–208.

- [13] A. Tyurin, *Analogues of Torelli's theorem for multidimensional vector bundles over an arbitrary algebraic curve*, Izv. Akad. Nauk SSSR Ser. Mat. **34** (1970), 338–365. English translation: Math. USSR – Izv. **4** (1970), 343–370.

CHENNAI MATHEMATICAL INSTITUTE, 92, G.N. CHETTY ROAD, CHENNAI 600017, INDIA

*E-mail address:* `balaji@cmi.ac.in`

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA ROAD, MUMBAI 400005, INDIA

*E-mail address:* `indranil@math.tifr.res.in`

THE INSTITUTE OF MATHEMATICAL SCIENCES, CIT CAMPUS, TARAMANI, CHENNAI 600113, INDIA

*E-mail address:* `dsn@imsc.res.in`

DEPARTMENT OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF LIVERPOOL, PEACH STREET, LIVERPOOL, L69 7ZL, ENGLAND

*E-mail address:* `newstead@liverpool.ac.uk`